

**Table 1. Cardiac Evaluation in Athletes with Prior COVID-19 Infection**

Clinical Scenario	Recommended Assessment	Comments
Athletes with <b>prior asymptomatic infection</b> as confirmed antibody to SARS-Coronavirus-2	<p><u>Focused Medical History and Physical Examination</u> to screen for findings newly emergent in the COVID-19 era.</p> <p><b>Consider 12-lead ECG*</b></p> <ul style="list-style-type: none"> <li>If ECG is abnormal or shows new repolarization changes compared to a prior ECG, then additional evaluation with at minimum an echocardiogram and exercise test is warranted in conjunction with a sports cardiologist.</li> </ul>	<ul style="list-style-type: none"> <li>Myopericarditis related to COVID-19 should be considered in patients with a history of new onset chest pain/pressure (even in the absence of fever and respiratory symptoms), palpitations, or exercise intolerance.</li> <li>Comprehensive clinical evaluation, regardless of ECG findings, is indicated in athletes with new onset cardiovascular symptoms or exercise intolerance.</li> </ul>
Athletes with a <b>history of mild illness (non-hospitalized)</b> related to confirmed or suspected COVID-19	<p><u>Focused Medical History and Physical Examination</u> to screen for persistent or new post-infectious findings following COVID-19 infection.</p> <p><b>Perform 12-lead ECG*</b></p> <ul style="list-style-type: none"> <li>If ECG is abnormal or shows new repolarization changes compared to a prior ECG, then additional individualized evaluation is warranted, including at minimum echocardiography and exercise testing, in conjunction with a sports cardiologist.</li> </ul>	<ul style="list-style-type: none"> <li><b>ECG findings</b> that may indicate viral-induced myocardial injury include: pathological Q waves, ST segment depression, (new) diffuse ST segment elevation, and T-wave inversion.</li> <li>Comprehensive clinical evaluation, regardless of ECG findings, is indicated in athletes with new onset cardiovascular symptoms or exercise intolerance.</li> </ul>
Athletes with a <b>history of moderate to severe illness (hospitalized)</b> related to confirmed or suspected COVID-19	<p><b>Comprehensive evaluation</b> prior to return to sport, in conjunction with a sports cardiologist, to include blood biomarker assessment (i.e. hs-Tn, NP), 12-lead ECG, echocardiography, exercise testing, and ambulatory rhythm monitoring.</p>	<ul style="list-style-type: none"> <li><b>Myocardial injury is more likely in patients with a more severe disease</b> course, and normal cardiac function and exercise tolerance should be established prior to a return to exercise.</li> <li>Cardiac MRI may be considered based on clinical suspicion of myocardial injury.**</li> </ul>

